

A Note on the Formulas for the Drazin Inverse of the Sum of Two Matrices and Its Applications

Xiaolan Qin^{*}, Ricai Luo

School of Mathematics and Statistics, Hechi University, Yizhou, China

Email address:

qinxiaolan18@163.com (Xiaolan Qin), luoricai@163.com (Rcai Luo)

^{*}Corresponding author

To cite this article:

Xiaolan Qin, Ricai Luo. A Note on the Formulas for the Drazin Inverse of the Sum of Two Matrices and Its Applications. *Pure and Applied Mathematics Journal*. Vol. 8, No. 3, 2019, pp. 54-71. doi: 10.11648/j.pamj.20190803.12

Received: July 6, 2019; Accepted: August 10, 2019; Published: August 30, 2019

Abstract: The Drazin inverse has applications in a number of areas such as control theory, Markov chains, singular differential and difference equations, and iterative methods in numerical linear algebra. The study on representations for the Drazin inverse of block matrices stems essentially from finding the general expressions for the solutions to singular systems of differential equations, and then stimulated by a problem formulated by Campbell. In 1983, Campbell (Campbell et al. (1976)) established an explicit representation for the Drazin inverse of a 2×2 block matrix M in terms of the blocks of the partition, where the blocks A and D are assumed to be square matrices. Special cases of the problems have been studied. In 2009, Chunyuan Deng and Yimin Wei found an explicit representation for the Drazin inverse of an anti-triangular matrix M , where A and BC are generalized Drazin invertible, if $A^\pi AB=0$ and $BC(I-A^\pi)=0$. Afterwards, several authors have investigated this problem under some limited conditions on the blocks of M . In particular, a representation of the Drazin inverse of M , denoted by M^d . In this paper, we consider the Drazin inverse of a sum of two matrices and we derive additive formulas under the conditions of $ABA^\pi=0$ and $BA^\pi=0$ respectively. Precisely, for a block matrix M , we give a new representation of M^d under some conditions that $AB=0$ and $DCA^\pi=0$. Moreover, some particular cases of this result related to the Drazin inverse of block matrices are also considered.

Keywords: Drazin Inverse, Block Matrices, Drazin Index

1. Introduction

Let A be a square complex matrix. We denote by $R(A)$, $N(A)$ and $\text{rank}(A)$, the range, the null space and the rank of matrix A , respectively. In addition, the smallest integer k is called the Drazin index of A such that $\text{rank}(A^{k+1}) = \text{rank}(A^k)$. Let $\mathbb{C}^{n \times n}$ be the set of all the $n \times n$ matrices over the complex field, for every matrix $A \in \mathbb{C}^{n \times n}$, such that $\text{index}(A) = k$, there exists a unique matrix $A^d \in \mathbb{C}^{n \times n}$, which satisfies the relations:

$$A^d A A^d = A^d, A^d A = A A^d, A^{k+1} A^d = A^k.$$

The matrix A^d is called the Drazin inverse of A [1, 2]. The case $\text{index}(A) = 0$ is valid if and only if A is nonsingular, so A^d reduces to A^{-1} . By $A^\pi = I - A A^d$, we

denote the projection on $N(A^k)$ along $R(A^k)$. If the lower limit of a sum is greater than its upper limit, we always define

the sum to be 0. For example, the sum $\sum_{l=1}^{k-2} * = 0$ for $k \leq 2$. We

agree that $A^0 = I$, for any matrix A .

The study on representations for the Drazin inverse of block matrices essentially originated from finding the general expressions for the solutions to singular systems of differential equations [3-5], and then stimulated by a problem formulated by Campbell [3]: establish an explicit representation for the Drazin inverse of 2×2 block matrices

$$M = \begin{bmatrix} A & B \\ C & D \end{bmatrix}, \quad (1)$$

in terms of the blocks of the partition, where the blocks A and D are assumed to be square matrices. For a deeper discussion

of applications of the Drazin inverse of a 2×2 block matrix, we refer the reader to [5, 6]. Meyer and Rose [7], and independently Hartwig and Shoaf [8], first gave the formulas for block triangular matrices. Several authors have investigated this problem and they were able to find some partial answers (imposing some conditions on the blocks of M). Here, we list some cases of Drazin inverse of block matrix M :

- (1) $AB = 0$ and $DC = 0$. See [9].
- (2) $BC = 0$ and $BD = 0$. See [10].
- (3) $BC = 0$ and $DC = 0$. See [11].
- (4) $AA^\pi B = 0$, $BCAA^d = 0$ and $DC = 0$. See [12].
- (5) $BCA = 0, BCB = 0, DCA = 0$ and $DCB = 0$. See [13].
- (6) $BCA = 0$, $CBD = 0$, $A(BC)^\pi = 0$ and $D(CB)^\pi = 0$.

See [14].

The motivation for this article is [15]. In the paper, the authors considered some conditions on $a, b \in \Lambda$ (Let Λ be a complex Banach algebra with the unit 1) that allowed them to express $(a + b)^d$ in terms of a, ad, b, bd . In this paper, we consider the Drazin inverse of a sum of two matrices and we derive additive formulas under the conditions of $ABA^\pi = 0$ and $BA^\pi = 0$ respectively. As an application we give some new representations for the Drazin inverse of a block matrix.

2. A New Additive Result for the Drazin Inverse of Matrices

First, we will state some auxiliary lemmas.

Lemma 2.1 [16] Let $A, B \in \mathbb{C}^{n \times n}$. If $AB = 0$ and A is nilpotent, then

$$(A + B)^d = \sum_{i=0}^{r-1} (B^d)^{i+1} A^i, \quad \text{ind}(A) = r$$

Lemma 2.2 [7] Let M_1 and M_2 be complex matrices of the form

$$M_1 = \begin{bmatrix} A & 0 \\ C & B \end{bmatrix}, \quad M_2 = \begin{bmatrix} B & C \\ 0 & A \end{bmatrix},$$

where A and B are complex square matrices. Let $r = \text{ind}(A)$ and $s = \text{ind}(B)$. Then $\max\{r, s\} \leq \text{ind}(M_i) \leq r + s$, for $i = 1, 2$, and

$$M_1^d = \begin{bmatrix} A^d & 0 \\ S & B^d \end{bmatrix}, \quad M_2^d = \begin{bmatrix} B^d & S \\ 0 & A^d \end{bmatrix},$$

$$0 = ABA^\pi = P \begin{bmatrix} A_1 & 0 \\ 0 & A_2 \end{bmatrix} \begin{bmatrix} B_1 & B_2 \\ B_3 & B_4 \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 0 & I \end{bmatrix} P^{-1} = P \begin{bmatrix} 0 & A_1 B_2 \\ 0 & A_2 B_4 \end{bmatrix} P^{-1}.$$

Therefore $A_1 B_2 = 0$ and $A_2 B_4 = 0$. The nonsingular of A_1 leads to $B_2 = 0$. Hence,

where

$$S = (B^d)^2 \sum_{i=0}^{r-1} (B^d)^i C A^i A^\pi + B^\pi \sum_{i=0}^{s-1} B^i C (A^d)^i (A^d)^2 - B^d C A^d.$$

The following theorem, we obtain the same expression for the Drazin inverse $(A + B)^d$ as in [15, Theorem 2.3] for the Generalized Drazin inverse in a Banach Algebra.

Theorem 2.1. Let $A, B \in \mathbb{C}^{n \times n}$. If $ABA^\pi = 0$, we have

$$\begin{aligned} (A + B)^d &= W^d + \sum_{i=0}^{r-1} (B^d)^{i+1} A^i A^\pi - \sum_{i=0}^{r-1} (B^d)^{i+1} A^i A^\pi B W^d \\ &\quad + \sum_{j=0}^{k-1} \left(\sum_{i=0}^{r-1} (B^d)^{i+j+2} A^i \right) A^\pi B W^j W^\pi \\ &\quad + B^\pi \sum_{j=0}^{k-1} (A + B)^j A^\pi B (W^d)^{j+2} \\ &\quad - \sum_{j=0}^{k-1} \left(\sum_{i=0}^{r-1} (B^d)^{i+1} A^{i+1} \right) (A + B)^j A^\pi B (W^d)^{j+2}, \end{aligned}$$

where $W = AA^d(A + B)$ and $\text{ind}(A) = r$, $\text{ind}(B) = t$, $\max\{\text{ind}(A), \text{ind}(B)\} \leq k \leq \text{ind}(A) + \text{ind}(B)$, $\text{ind}(W) \leq \text{ind}(A) + \text{ind}(B)$.

Proof. If we represent

$$A \text{ as } A = P \begin{bmatrix} A_1 & 0 \\ 0 & A_2 \end{bmatrix} P^{-1},$$

where P and A_1 are nonsingular and A_2 is nilpotent, then

$$A^d = P \begin{bmatrix} A_1^{-1} & 0 \\ 0 & 0 \end{bmatrix} P^{-1}$$

and $\text{ind}(A) = \text{ind}(A_1)$. Let us write

$$B = P \begin{bmatrix} B_1 & B_2 \\ B_3 & B_4 \end{bmatrix} P^{-1},$$

where $B_1 \in \mathbb{C}^{r \times r}$ being r the size of A_1 .

From $ABA^\pi = 0$, we have

$$B = P \begin{bmatrix} B_1 & 0 \\ B_3 & B_4 \end{bmatrix} P^{-1}, A + B = P \begin{bmatrix} A_1 + B_1 & 0 \\ B_3 & A_2 + B_4 \end{bmatrix} P^{-1}. \quad (2)$$

Observe that

$$W = AA^d(A+B) = P \begin{bmatrix} A_1 & 0 \\ 0 & A_2 \end{bmatrix} \begin{bmatrix} A_1^{-1} & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} A_1 + B_1 & 0 \\ B_3 & A_2 \end{bmatrix} P^{-1} = P \begin{bmatrix} A_1 + B_1 & 0 \\ 0 & 0 \end{bmatrix} P^{-1},$$

which leads to

$$W^d = P \begin{bmatrix} (A_1 + B_1)^d & 0 \\ 0 & 0 \end{bmatrix} P^{-1}.$$

Since A_2 is nilpotent and $A_2 B_4 = 0$, Lemma 2.1 yields

$$(A_2 + B_4)^d = \sum_{i=0}^{r-1} (B_4^d)^{i+1} A_2^i. \quad (3)$$

Thus, by lemma 2.2, we obtain

$$(A+B)^d = P \begin{bmatrix} (A_1 + B_1)^d & 0 \\ S & (A_2 + B_4)^d \end{bmatrix} P^{-1},$$

and

$$S = \sum_{j=0}^{p-1} ((A_2 + B_4)^d)^{j+2} B_3 (A_1 + B_1)^j (A_1 + B_1)^\pi + \sum_{j=0}^{q-1} (A_2 + B_4)^\pi (A_2 + B_4)^j B_3 ((A_1 + B_1)^d)^{j+2} - (A_2 + B_4)^d B_3 (A_1 + B_1)^d, \quad (4)$$

where $p = \text{ind}(A_1 + B_1)$ and $q = \text{ind}(A_2 + B_2)$. Observe that for any $X \in \mathbb{C}^{m \times m}$, one has $\text{ind}(X) \leq m$ and if $k \geq \text{ind}(X)$, then $X^k X^\pi = 0$ (both affirmations can be proved by means of the Jordan canonical form of X). Thus, in (4) the upper limits of the summations can be replaced by simply n .

We have (we will write with an asterisk any entry whose exactly expression is not necessary)

$$(B^d)^{i+1} A^i A^\pi = P \begin{bmatrix} (B_1^d)^{i+1} & 0 \\ * & (B_4^d)^{i+1} \end{bmatrix} \begin{bmatrix} A_1^i & 0 \\ 0 & A_2^i \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 0 & I_{n-r} \end{bmatrix} P^{-1} = P \begin{bmatrix} 0 & 0 \\ 0 & (B_4^d)^{i+1} A_2^i \end{bmatrix} P^{-1}.$$

Hence, we get

$$\sum_{i=0}^{r-1} (B^d)^{i+1} A^i A^\pi = \sum_{i=0}^{r-1} (B^d)^{i+1} A^i A^\pi = P \begin{bmatrix} 0 & 0 \\ 0 & \sum_{i=0}^{r-1} (B_4^d)^{i+1} A_2^i \end{bmatrix} P^{-1} = P \begin{bmatrix} 0 & 0 \\ 0 & (A_2 + B_4)^d \end{bmatrix} P^{-1}.$$

Therefore,

$$\sum_{i=0}^{r-1} (B^d)^{i+1} A^i A^\pi B W^d = P \begin{bmatrix} 0 & 0 \\ 0 & (A_2 + B_4)^d \end{bmatrix} \begin{bmatrix} B_1 & 0 \\ B_3 & B_4 \end{bmatrix} \begin{bmatrix} (A_1 + B_1)^d & 0 \\ 0 & 0 \end{bmatrix} P^{-1} = P \begin{bmatrix} 0 & 0 \\ (A_2 + B_4)^d B_3 (A_1 + B_1)^d & 0 \end{bmatrix} P^{-1}.$$

In a similar way, we get for any $j \in \mathbb{N}$

$$\left(\sum_{i=0}^{r-1} (B^d)^{i+1} A^i A^\pi \right)^{j+2} B W^j W^\pi = P \begin{bmatrix} 0 & 0 \\ \left[(A_2 + B_4)^d \right]^{j+2} B_3 (A_1 + B_1)^j (A_1 + B_1)^\pi & 0 \end{bmatrix} P^{-1}.$$

Now, we will find an expression for $(A_2 + B_4)^\pi$. To this end, we use $A_2 B_4 = 0$ and (3). Observe that

$$A_2 B_4^d = A_2 B_4 (B_4^d)^2 = 0.$$

$$\begin{aligned} (A_2 + B_4)^\pi &= I_{n-r} - (A_2 + B_4)(A_2 + B_4)^d \\ &= I_{n-r} - (A_2 + B_4) \left[B_4^d + (B_4^d)^2 A_2 + (B_4^d)^3 A_2^2 + \cdots + (B_4^d)^r A_2^{r-1} \right] \\ &= I_{n-r} - \left[B_4 B_4^d + B_4 (B_4^d)^2 A_2 + B_4 (B_4^d)^3 A_2^2 + \cdots + B_4 (B_4^d)^r A_2^{r-1} \right] \\ &= B_4^\pi - \left[B_4^d A_2 + (B_4^d)^2 A_2^2 + \cdots + (B_4^d)^{r-1} A_2^{r-1} \right], \end{aligned}$$

and so,

$$\begin{aligned} \sum_{i=0}^{r-1} (B^d)^{i+1} A^{i+1} A^\pi &= B^d A A^\pi + (B^d)^2 A^2 A^\pi + \cdots + (B^d)^{r-1} A^{r-1} A^\pi \\ &= P \left\{ \begin{bmatrix} 0 & 0 \\ 0 & B_4^d A_2 \end{bmatrix} + \cdots + \begin{bmatrix} 0 & 0 \\ 0 & (B_4^d)^{r-1} A_2^{r-1} \end{bmatrix} \right\} P^{-1} \\ &= P \begin{bmatrix} 0 & 0 \\ 0 & B_4^\pi - (A_2 + B_4)^\pi \end{bmatrix} P^{-1}. \end{aligned}$$

In addition,

$$\begin{aligned} B^\pi &= I - B B^d \\ &= P \left\{ \begin{bmatrix} I_r & 0 \\ 0 & I_{n-r} \end{bmatrix} - \begin{bmatrix} B_1 & 0 \\ B_3 & B_4 \end{bmatrix} \begin{bmatrix} B_1^d & 0 \\ * & B_4^d \end{bmatrix} \right\} P^{-1} \\ &= P \begin{bmatrix} I_r & 0 \\ * & B_4^\pi \end{bmatrix} P^{-1}, \end{aligned}$$

which implies

$$B^\pi A^\pi = P \begin{bmatrix} 0 & 0 \\ 0 & B_4^\pi \end{bmatrix} P^{-1}.$$

Thus,

$$\begin{aligned} P \begin{bmatrix} 0 & 0 \\ 0 & (A_2 + B_4)^\pi \end{bmatrix} P^{-1} &= P \begin{bmatrix} 0 & 0 \\ 0 & B_4^\pi \end{bmatrix} P^{-1} - \sum_{i=0}^{r-1} (B^d)^{i+1} A^{i+1} A^\pi \\ &= B^\pi A^\pi - \sum_{i=0}^{r-1} (B^d)^{i+1} A^{i+1} A^\pi \\ &= (B^\pi - \sum_{i=0}^{r-1} (B^d)^{i+1} A^{i+1}) A^\pi. \end{aligned}$$

Hence for $j \in \mathbb{N}$,

$$\begin{aligned} P \begin{bmatrix} 0 & 0 \\ 0 & (A_2 + B_4)^\pi (A_2 + B_4)^j \end{bmatrix} P^{-1} &= P \begin{bmatrix} 0 & 0 \\ 0 & (A_2 + B_4)^\pi \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 0 & (A_2 + B_4)^j \end{bmatrix} P^{-1} \\ &= \left[(B^\pi - \sum_{i=0}^{r-1} (B^d)^{i+1} A^{i+1}) A^\pi \right] \left[(A + B)^j A^\pi \right]. \end{aligned}$$

But, as it is easy from (2), one has

$$A^\pi (A + B)^j A^\pi = (A + B)^j A^\pi.$$

Therefore,

$$\begin{aligned}
(B^\pi - \sum_{i=0}^{r-1} (B^d)^{i+1} A^{i+1})(A+B)^j A^\pi B(W^d)^{j+2} &= P \begin{bmatrix} 0 & 0 \\ 0 & (A_2 + B_4)^\pi (A_2 + B_4)^j \end{bmatrix} \begin{bmatrix} B_1 & 0 \\ B_3 & B_4 \end{bmatrix} \begin{bmatrix} [(A_1 + B_1)^d]^{j+2} & 0 \\ 0 & 0 \end{bmatrix} P^{-1} \\
&= P \begin{bmatrix} 0 & 0 \\ (A_2 + B_4)^\pi (A_2 + B_4)^j B_3 [(A_1 + B_1)^d]^{j+2} & 0 \end{bmatrix} P^{-1}.
\end{aligned}$$

Finally, let us observe that the expression

$$\left(\sum_{i=0}^{r-1} (B^d)^{i+1} A^i A^\pi \right)^{j+2},$$

in effect, since

$$[(A_2 + B_4)^d]^{j+2} = \sum_{i=0}^{r-1} (B_4^d)^{i+j+2} A_2^i,$$

we have that

$$\left(\sum_{i=0}^{r-1} (B^d)^{i+1} A^i A^\pi \right)^{j+2} = \sum_{i=0}^{r-1} (B^d)^{i+j+2} A^i A^\pi$$

The proof is finished.

The next result, we obtain the same expression for the Drazin inverse $(A+B)^d$ as in [15, Theorem 2.6] for the Generalized Drazin inverse in a Banach Algebra. Theorem 2.2. Let $A, B \in \mathbb{C}^{n \times n}$ and $\text{ind}(A) = r$. If $BA^\pi = 0$, we have

$$(A+B)^d = W^d + A^\pi \sum_{i=0}^{r-1} A^i B(W^d)^{i+2},$$

where $W = AA^d(A+B)$.

Proof. We can represent A , A^d and B as in Theorem 2.1. From $BA^\pi = 0$, we have

$$0 = BA^\pi = P \begin{bmatrix} B_1 & B_2 \\ B_3 & B_4 \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 0 & I \end{bmatrix} P^{-1} = P \begin{bmatrix} 0 & B_2 \\ 0 & B_4 \end{bmatrix} P^{-1}.$$

Therefore $B_2 = B_4 = 0$. Hence,

$$B = P \begin{bmatrix} B_1 & 0 \\ B_3 & 0 \end{bmatrix} P^{-1}, \quad A+B = P \begin{bmatrix} A_1 + B_1 & 0 \\ B_3 & A_2 \end{bmatrix} P^{-1}.$$

Observe that

$$W = AA^d(A+B) = P \begin{bmatrix} A_1 & 0 \\ 0 & A_2 \end{bmatrix} \begin{bmatrix} A_1^{-1} & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} A_1 + B_1 & 0 \\ B_3 & A_2 \end{bmatrix} P^{-1} = P \begin{bmatrix} A_1 + B_1 & 0 \\ 0 & 0 \end{bmatrix} P^{-1},$$

which leads to

$$W^d = P \begin{bmatrix} (A_1 + B_1)^d & 0 \\ 0 & 0 \end{bmatrix} P^{-1}.$$

Thus, by Lemma 2.2, we obtain

$$(A+B)^d = P \begin{bmatrix} (A_1 + B_1)^d & 0 \\ S & 0 \end{bmatrix} P^{-1},$$

where

$$S = \sum_{i=0}^{p-1} A_2^i B_3 \left[(A_1 + B_1)^d \right]^{i+2},$$

and $p = \text{ind}(A_2)$.

Also we have

$$A^\pi A^i B (W^d)^{i+2} = P \begin{bmatrix} 0 & 0 \\ 0 & I_{n-r} \end{bmatrix} \begin{bmatrix} A_1^i & 0 \\ 0 & A_2^i \end{bmatrix} \begin{bmatrix} B_1 & 0 \\ B_3 & 0 \end{bmatrix} \begin{bmatrix} ((A_1 + B_1)^d)^{i+2} & 0 \\ 0 & 0 \end{bmatrix} = P \begin{bmatrix} 0 & 0 \\ A_2^i B_3 \left[(A_1 + B_1)^d \right]^{i+2} & 0 \end{bmatrix} P^{-1}.$$

Hence the Theorem follows.

3. Some Results on the Drazin Inverse of 2×2 Block Matrices

In this section we shall apply Theorem 2.1 and Theorem 2.2 to obtain some formulas for M^d under some conditions when M is a 2×2 block matrix written as in (1). We assume that $\text{ind}(A) = r$, $\text{ind}(D) = s$, $\text{ind}(BC) = t$, $\text{ind}(CB) = l$, and $\max\{\text{ind}(A), \text{ind}(D), \text{ind}(BC), \text{ind}(CB)\} \leq k \leq \text{ind}(A) + \text{ind}(D) + \text{ind}(BC) + \text{ind}(CB)$.

Next we will state some auxiliary lemmas.

Lemma 3.1 Let $A \in \mathbb{C}^{m \times n}$, $B \in \mathbb{C}^{n \times m}$.

$A((BA)^d)^i = ((AB)^d)^i A$ for every integer $i \geq 1$, and $B(AB)^\pi = (BA)^\pi B$. Moreover, $\text{ind}(BA) - 1 \leq \text{ind}(AB) \leq \text{ind}(BA) + 1$.

Proof. As in the proof of [5, Theorem 7.8.4], we can obtain $(AB)^d = A((BA)^d)^2 B$. The results follow.

Lemma 3.2 Let $X \in \mathbb{C}^{n \times n}$. Then $(XX^\pi)^d = 0$, $(X^2 X^d)^d = X^d$, $(X^2 X^d)^\pi = X^\pi$, and $\text{ind}(XX^\pi) = \text{ind}(X)$, $\text{ind}(X^2 X^d) = 1$.

Proof. The Jordan canonical form of X permits write $X = S(C \oplus N)S^{-1}$, where S and C are nonsingular, and N is nilpotent. Evidently, $X^d = S(C^{-1} \oplus 0)S^{-1}$. Now, it is evident $X^2 X^d = S(C \oplus 0)S^{-1}$ and $XX^\pi = S(0 \oplus N)S^{-1}$, which leads to the affirmations of this lemma.

Lemma 3.3 [10] If M is matrix of a form (1) such that $BC = 0$ and $BD = 0$, then

$$M^d = \begin{bmatrix} A^d & (A^d)^2 B \\ \Sigma_0 & D^d + \Sigma_1 B \end{bmatrix},$$

where

$$\Sigma_n = \sum_{i=0}^{r-1} (D^d)^{i+n+2} C A^i A^\pi + D^\pi \sum_{i=0}^{s-1} D^i C (A^d)^{i+n+2} - \sum_{i=0}^n (D^d)^{i+1} C (A^d)^{n-i+1}, n \geq 0. \quad (5)$$

Lemma 3.4 [17] Let M be a matrix of a the form (1) with $A = 0$ and $D = 0$, then

$$M^d = \begin{bmatrix} 0 & B(CB)^d \\ (CB)C^d & 0 \end{bmatrix}.$$

Furthermore, If $\text{ind}(BC) = l$, then $\text{ind}(M) \leq 2l + 1$.

Lemma 3.5 [18] If M is matrix of a form (1) such that $AB = 0$ and $D = 0$, then

$$M^d = \begin{bmatrix} XA & (BC)^d B \\ CX & 0 \end{bmatrix},$$

where

$$X = (BC)^\pi \sum_{i=0}^{t-1} (BC)^i (A^d)^{2i+2} + \sum_{i=0}^{\left[\frac{r}{2}\right]-1} \left[(BC)^d \right]^{i+1} A^{2i} A^\pi.$$

Using Theorem 2.1 and the previous lemmas, we get the following results.

Theorem 3.1 Let M be given by (1). If $AB = 0$ and $DCA^\pi = 0$, then

$$M^d = \begin{bmatrix} A^d + 2XA - RD^\pi CA^d & RD^\pi + (BC)^d BD^\pi \\ (CB)^\pi Z_0 + CX - SD^\pi CA^d + C(BC)^d (A^\pi + XA^2) & (CB)^\pi D^d + SD^\pi \end{bmatrix} \\ + \begin{bmatrix} I - BCX & -RD \\ -CXA & (CB)^\pi - SD \end{bmatrix} \sum_{j=0}^{k-1} \begin{bmatrix} A & B \\ C & D \end{bmatrix}^j \begin{bmatrix} BZ_{j+1} & B(D^d)^{j+2} \\ D^\pi C(A^d)^{j+2} & 0 \end{bmatrix},$$

where

$$X = \sum_{n=0}^{\left[\frac{r}{2}\right]-1} \left[(BC)^d \right]^{i+1} A^{2i} A^\pi, Z_n = - \sum_{i=0}^n (D^d)^{i+1} C(A^d)^{n-i+1}, \quad (6)$$

$$R = \sum_{i=0}^{\left[\frac{s}{2}\right]-1} \left[(BC)^d \right]^{i+1} BD^{2i}, S = \sum_{i=0}^{\left[\frac{s}{2}\right]-1} \left[(CB)^d \right]^{i+1} D^{2i+1}. \quad (7)$$

Proof. We can split matrix M as $M = P + Q$, where

$$P = \begin{bmatrix} A^2 A^d & 0 \\ 0 & D \end{bmatrix}, Q = \begin{bmatrix} AA^\pi & B \\ C & 0 \end{bmatrix}, \\ P^d = \begin{bmatrix} A^d & 0 \\ 0 & D^d \end{bmatrix}, P^\pi = \begin{bmatrix} A^\pi & 0 \\ 0 & D^\pi \end{bmatrix}.$$

From $AB = 0$ and $DCA^\pi = 0$, we have $PQP^\pi = 0$. Applying Theorem 2.1, we get

$$M^d = W^d + \sum_{i=0}^{p-1} (Q^d)^{i+1} P^i P^\pi - \sum_{i=0}^{p-1} (Q^d)^{i+1} P^i P^\pi Q W^d + \sum_{j=0}^{h-1} \left(\sum_{i=0}^{p-1} (Q^d)^{i+j+2} P^i \right) P^\pi Q W^j W^\pi + Q^\pi \sum_{j=0}^{h-1} (P + Q)^j P^\pi Q (W^d)^{j+2} \\ - \sum_{j=0}^{h-1} \sum_{i=0}^{p-1} (Q^d)^{i+1} P^{i+1} (P + Q)^j P^\pi Q (W^d)^{j+2} \quad (8)$$

where

$$\text{ind}(P) = p, \text{ind}(Q) = q, \text{ind}(Q) = q,$$

$$(\max\{\text{ind}(P), \text{ind}(Q)\} \leq h \leq \text{ind}(P) + \text{ind}(Q), \text{ind}(W) \leq \text{ind}(P) + \text{ind}(Q).$$

Now we consider the matrices mentioned in the above equation. Clearly, for every integer $i \geq 1$,

$$P^{2i} = \begin{bmatrix} A^{2i+1} A^d & 0 \\ 0 & D^{2i} \end{bmatrix}, P^{2i+1} = \begin{bmatrix} A^{2i+1} A^d & 0 \\ 0 & D^{2i+1} \end{bmatrix}.$$

Since $AB = 0$, we have $AA^\pi B = 0$ and matrix Q satisfies the conditions of Lemma 3.5, so we get

$$Q^d = \begin{bmatrix} XA & (BC)^d B \\ CX & 0 \end{bmatrix}, Q^\pi = \begin{bmatrix} I - BCX & 0 \\ -CXA & (CB)^\pi \end{bmatrix},$$

where X is defined in (6).

From $AB = 0$, we have $AX = 0$, $XX = (BC)^d X$,

$XB = (BC)^d B$, $BC(BC)^d X = X$ and $X(BC)^d = ((BC)^d)^2$. Then, by Lemma 3.1, for every integer $i \geq 1$,

$$(Q^d)^{2i} = \begin{bmatrix} [(BC)^d]^{i-1} X & 0 \\ C[(BC)^d]^i XA & [(CB)^d]^i \end{bmatrix},$$

$$(Q^d)^{2i+1} = \begin{bmatrix} [(BC)^d]^i XA & [(BC)^d]^{i+1} B \\ C[(BC)^d]^i X & 0 \end{bmatrix}.$$

From $AB = 0$, we have

$$W = PP^d(P+Q) = \begin{bmatrix} A^2 A^d & AA^d B \\ DD^d C & D^2 D^d \end{bmatrix} = \begin{bmatrix} A^2 A^d & 0 \\ DD^d C & D^2 D^d \end{bmatrix},$$

by Lemma 2.2 and Lemma 3.2, we obtain

$$W^d = \begin{bmatrix} A^d & 0 \\ Z_0 & D^d \end{bmatrix}, (W^d)^n = \begin{bmatrix} (A^d)^n & 0 \\ Z_{n-1} & (D^d)^n \end{bmatrix}, W^\pi = \begin{bmatrix} A^\pi & 0 \\ 0 & D^\pi \end{bmatrix}$$

where Z_n is defined in (6).

Since $XAA^\pi = XA$, we have, for $i = 0$,

$$\sum_{i=0}^{p-1} (Q^d)^{i+1} P^i P^\pi = Q^d P^\pi + (Q^d)^2 PP^\pi = \begin{bmatrix} XA & (BC)^d BD^\pi \\ CX & (CB)^d DD^\pi \end{bmatrix},$$

since $XAA^d = 0$, we have, for every integer $i \geq 1$,

$$\sum_{i=0}^{\left[\frac{p}{2}\right]-1} (Q^d)^{2i+1} P^{2i} = \sum_{i=0}^{\left[\frac{p}{2}\right]-1} \begin{bmatrix} [(BC)^d]^i XA^{2i+2} A^d & [(BC)^d]^{i+1} BD^{2i} \\ C[(BC)^d]^i XA^{2i+1} A^d & 0 \end{bmatrix} = \begin{bmatrix} 0 & \sum_{i=0}^{\left[\frac{s}{2}\right]-1} [(BC)^d]^{i+1} BD^{2i} \\ 0 & 0 \end{bmatrix},$$

and

$$\sum_{n=0}^{\left[\frac{p}{2}\right]-1} (Q^d)^{2i+2} P^{2i+1} = \sum_{i=0}^{\left[\frac{p}{2}\right]-1} \begin{bmatrix} [(BC)^d]^i XA^{2i+2} A^d & 0 \\ C[(BC)^d]^{i+1} XA^{2i+3} A^d & [(CB)^d]^{i+1} D^{2i+1} \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & \sum_{i=0}^{\left[\frac{s}{2}\right]-1} [(CB)^d]^{i+1} D^{2i+1} \end{bmatrix},$$

And, similarly

$$\sum_{i=0}^{\left[\frac{p}{2}\right]-1} (Q^d)^{2i+1} P^{2i+1} = \begin{bmatrix} 0 & \sum_{i=0}^{\left[\frac{s}{2}\right]-1} [(BC)^d]^{i+1} BD^{2i+1} \\ 0 & 0 \end{bmatrix} \quad (9)$$

and

$$\sum_{i=0}^{\left[\frac{p}{2}\right]-1} (Q^d)^{2i+2} P^{2i+2} = \begin{bmatrix} 0 & 0 \\ 0 & \sum_{i=0}^{\left[\frac{s}{2}\right]-1} [(CB)^d]^{i+1} BD^{2i+2} \end{bmatrix} \quad (10)$$

Hence the first sum in (8) is,

$$\sum_{i=0}^{p-1} (Q^d)^{i+1} P^i P^\pi = \left[\sum_{i=0}^{\left[\frac{p}{2}\right]-1} (Q^d)^{2i+1} P^{2i} + \sum_{i=0}^{\left[\frac{p}{2}\right]-1} (Q^d)^{2i+2} P^{2i+1} \right] P^\pi = \begin{bmatrix} XA & \sum_{i=0}^{\left[\frac{s}{2}\right]-1} [(BC)^d]^{i+1} BD^{2i} D^\pi \\ CX & \sum_{i=0}^{\left[\frac{s}{2}\right]-1} [(CB)^d]^{i+1} D^{2i+1} D^\pi \end{bmatrix} = \begin{bmatrix} XA & RD^\pi \\ CX & SD^\pi \end{bmatrix},$$

where R and S are defined in (7).

Next consider the second sum in (8).

Note that

$$QW^d = \begin{bmatrix} AA^\pi & B \\ C & 0 \end{bmatrix} \begin{bmatrix} A^d & 0 \\ Z_0 & D^d \end{bmatrix} = \begin{bmatrix} BZ_0 & BD^d \\ CA^d & 0 \end{bmatrix}.$$

Since $XB = (BC)^d B$ and $AB = 0$, using Lemma 3.1, we get

$$\sum_{i=0}^{p-1} (Q^d)^{i+1} P^i P^\pi QW^d = \begin{bmatrix} XA & RD^\pi \\ CX & SD^\pi \end{bmatrix} \begin{bmatrix} BZ_0 & BD^d \\ CA^d & 0 \end{bmatrix} = \begin{bmatrix} RD^\pi CA^d & 0 \\ CB(CB)^d Z_0 + SD^\pi CA^d & CB(CB)^d D^d \end{bmatrix}.$$

Since $DCA^\pi = 0$, we have

$$PP^\pi QW^\pi = \begin{bmatrix} 0 & 0 \\ 0 & DD^\pi \end{bmatrix} \begin{bmatrix} AA^\pi & B \\ C & 0 \end{bmatrix} \begin{bmatrix} A^\pi & 0 \\ 0 & D^\pi \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ DD^\pi CA^\pi & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix},$$

and

$$WW^\pi = \begin{bmatrix} A^2 A^d & 0 \\ DD^d C & D^2 D^d \end{bmatrix} \begin{bmatrix} A^\pi & 0 \\ 0 & D^\pi \end{bmatrix} = \begin{bmatrix} A^2 A^d A^\pi & 0 \\ DD^d CA^\pi & D^2 D^d D^\pi \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}.$$

Since $XB = (BC)^d B$, $AB = 0$ and $DCA^\pi = 0$, we can prove

$$\begin{aligned} \sum_{j=0}^{h-1} \left(\sum_{i=0}^{p-1} (Q^d)^{i+j+2} P^i \right) P^\pi QW^j W^\pi &= (Q^d)^2 P^\pi QW^\pi = \begin{bmatrix} X & 0 \\ C(BC)^d XA & (CB)^d \end{bmatrix} \begin{bmatrix} AA^\pi & A^\pi BD^\pi \\ D^\pi CA^\pi & 0 \end{bmatrix} \\ &= \begin{bmatrix} XAA^\pi & XBD^\pi \\ C(BC)^d XA^2 A^\pi + (CB)^d D^\pi CA^\pi & C(BC)^d XAA^\pi BD^\pi \end{bmatrix} = \begin{bmatrix} XA & (BC)^d BD^\pi \\ C(BC)^d XA^2 + (CB)^d CA^\pi & 0 \end{bmatrix}. \end{aligned}$$

Observe that (9) and (10) yield

$$\sum_{i=0}^{p-1} (Q^d)^{i+1} P^{i+1} = \sum_{i=0}^{\left[\frac{p}{2}\right]-1} (Q^d)^{2i+1} P^{2i+1} + \sum_{i=0}^{\left[\frac{p}{2}\right]-1} (Q^d)^{2i+2} P^{2i+2} = \begin{bmatrix} 0 & RD \\ 0 & SD \end{bmatrix},$$

So we get

$$Q^\pi \sum_{j=0}^{h-1} (P+Q)^j P^\pi Q(W^d)^{j+2} - \sum_{j=0}^{h-1} \sum_{i=0}^{p-1} (Q^d)^{i+1} P^{i+1} (P+Q)^j P^\pi Q(W^d)^{j+2},$$

$$= \begin{bmatrix} I - BCX & -RD \\ -CXA & (CB)^\pi - SD \end{bmatrix} \sum_{j=0}^{k-1} \begin{bmatrix} A & B \\ C & D \end{bmatrix}^j \begin{bmatrix} BZ_{j+1} & B(D^d)^{j+2} \\ D^\pi C(A^d)^{j+2} & 0 \end{bmatrix}.$$

The proof is finished

If we assume that $ABD^\pi = 0$ and $DD^dC = 0$ instead of $AB = 0$, we will get another expression for M^d .

Theorem 3.2 Let M be given by (1). If $ABD^\pi = 0$, $DCA^\pi = 0$ and $DD^dC = 0$, then

$$M^d = \begin{bmatrix} A^d + XA + (BC)^\pi L - BY(CA^d + DS) \\ CX - YDCA^d + C(BC)^d A^\pi + (CB)^\pi S - YD^2S \\ Z_1 - XA(BD^d + AN) + B(CB)^d D^\pi + (BC)^\pi N + BY(I - CZ_1 - DR) \\ D^d - CX(BD^d + AN) + (CB)^\pi R + YD(I - CZ_1 - DR) \end{bmatrix}$$

where

$$X = \sum_{n=0}^{\left[\frac{r}{2}\right]-1} [(BC)^d]^{n+1} A^{2n} A^\pi, Y = \sum_{n=0}^{\left[\frac{s}{2}\right]-1} [(CB)^d]^{n+1} D^{2n} D^\pi, Z_n = -\sum_{i=0}^{n-1} (A^d)^{i+1} B(D^d)^{n-i},$$

$$L = \sum_{n=1}^{k-1} B \left(\sum_{i=0}^{n-1} (CB)^i D^{2n-2i-1} C(A^d)^{2n+2} + \sum_{i=0}^n (CB)^i D^{2n-2i} C(A^d)^{2n+3} \right),$$

$$N = \sum_{n=1}^{k-1} B \left(\sum_{i=0}^{n-1} (CB)^i D^{2n-2i-1} CZ_{2n+2} + \sum_{i=0}^n (CB)^i D^{2n-2i} CZ_{2n+3} \sum_{i=0}^n (BC)^i A^{2n-2i} A^\pi (B + ABD^d) (D^d)^{2n+2} \right), \quad (11)$$

$$S = \sum_{n=1}^{k-1} \left(\sum_{i=0}^n (CB)^i D^{2n-2i} (C + DCA^d) (A^d)^{2n+2} \right),$$

$$R = \sum_{n=1}^{k-1} \left(\sum_{i=0}^n (CB)^i D^{2n-2i} (CZ_{2n+2} + DCZ_{2n+3}) + \sum_{i=0}^{n-1} (CB)^i CA^{2n-2i-1} A^\pi B(D^d)^{2n+2} + \sum_{i=0}^n (CB)^i CA^{2n-2i} A^\pi B(D^d)^{2n+3} \right)$$

Proof. We can split matrix M as $M = P + Q$, where

$$P = \begin{bmatrix} A & 0 \\ 0 & D \end{bmatrix}, Q = \begin{bmatrix} 0 & B \\ C & 0 \end{bmatrix}, P^d = \begin{bmatrix} A^d & 0 \\ 0 & D^d \end{bmatrix}, P^\pi = \begin{bmatrix} A^\pi & 0 \\ 0 & D^\pi \end{bmatrix}. \quad (12)$$

From lemma 3.4, we have

$$Q^d = \begin{bmatrix} 0 & B(CB)^d \\ (CB)^d C & 0 \end{bmatrix}, Q^\pi = \begin{bmatrix} (BC)^\pi & 0 \\ 0 & (CB)^\pi \end{bmatrix}. \quad (13)$$

From $ABD^\pi = 0$ and After the course, there were significant differences in the scores of heart sounds, lung sounds, mixed sound and total score between the two groups ($P < 0.05$), but there was no significant difference in the scores of bowl sound ($P > 0.05$).

we have $PQP^\pi = 0$. Applying Theorem 2.1, we get

$$\begin{aligned}
M^d &= W^d + \sum_{i=0}^{p-1} (Q^d)^{i+1} P^i P^\pi - \sum_{i=0}^{p-1} (Q^d)^{i+1} P^i P^\pi Q W^d \\
&+ \sum_{j=0}^{h-1} \left(\sum_{i=0}^{p-1} (Q^d)^{i+j+2} P^i \right) P^\pi Q W^j W^\pi + Q^\pi \sum_{j=0}^{h-1} (P+Q)^j P^\pi Q (W^d)^{j+2} - \sum_{j=0}^{h-1} \sum_{i=0}^{p-1} (Q^d)^{i+1} P^{i+1} (P+Q)^j P^\pi Q (W^d)^{j+2},
\end{aligned}$$

where

$$\text{ind}(P) = p, \text{ind}(Q) = q, \text{ind}(Q) = q, \max\{\text{ind}(P), \text{ind}(Q)\} \leq h \leq \text{ind}(P) + \text{ind}(Q) \text{ and } \text{ind}(W) \leq \text{ind}(P) + \text{ind}(Q),$$

From $DD^dC = 0$, we obtain

$$W = PP^d(P+Q) = \begin{bmatrix} A^2 A^d & AA^d B \\ DD^d C & D^2 D^d \end{bmatrix}. \quad (14)$$

By Lemma 2.2 and Lemma 3.2, we have

$$W^d = \begin{bmatrix} A^d & Z_1 \\ 0 & D^d \end{bmatrix}, (W^d)^n = \begin{bmatrix} (A^d)^n & Z_n \\ 0 & (D^d)^n \end{bmatrix}, W^\pi = \begin{bmatrix} A^\pi & 0 \\ 0 & D^\pi \end{bmatrix}, \quad (15)$$

where Z_n is defined in (11).

Clearly, for every integer $n \geq 1$,

$$(Q^d)^{2n} = \begin{bmatrix} ((BC)^d)^n & 0 \\ 0 & ((CB)^d)^n \end{bmatrix}, (Q^d)^{2n+1} = \begin{bmatrix} 0 & B((CB)^d)^{n+1} \\ ((CB)^d)^{n+1} C & 0 \end{bmatrix}.$$

Hence

$$\begin{aligned}
\sum_{i=0}^{p-1} (Q^d)^{i+1} P^i P^\pi &= \left[\sum_{i=0}^{\left[\frac{p}{2}\right]-1} (Q^d)^{2i+1} P^{2i} + \sum_{i=0}^{\left[\frac{p}{2}\right]-1} (Q^d)^{2i+2} P^{2i+1} \right] P^\pi \\
&= \begin{bmatrix} \sum_{n=0}^{\left[\frac{r}{2}\right]-1} ((BC)^d)^{i+1} A^{2i+1} A^\pi & \sum_{n=0}^{\left[\frac{s}{2}\right]-1} B((CB)^d)^{i+1} D^{2i} D^\pi \\ \sum_{n=0}^{\left[\frac{r}{2}\right]-1} C((BC)^d)^{i+1} A^{2i} A^\pi & \sum_{n=0}^{\left[\frac{s}{2}\right]-1} ((CB)^d)^{i+1} D^{2i+1} D^\pi \end{bmatrix} \\
&= \begin{bmatrix} XA & BY \\ CX & YD \end{bmatrix},
\end{aligned} \quad (16)$$

where X and Y are defined in (11).

$$QW^d = \begin{bmatrix} 0 & BD^d \\ CA^d & CZ_1 \end{bmatrix}. \quad (17)$$

By (16) and (17), we get

$$\sum_{i=0}^{p-1} (Q^d)^{i+1} P^i P^\pi QW^d = \begin{bmatrix} XA & BY \\ CX & YD \end{bmatrix} \begin{bmatrix} 0 & BD^d \\ CA^d & CZ_1 \end{bmatrix} = \begin{bmatrix} BYCA^d & XABD^d + BYCZ_1 \\ YDCA^d & CXBD^d + YDCZ_1 \end{bmatrix}.$$

Since $ABD^\pi = 0$, we have

$$WW^\pi = \begin{bmatrix} A^2 A^d & AA^d B \\ 0 & D^2 D^d \end{bmatrix} \begin{bmatrix} A^\pi & 0 \\ 0 & D^\pi \end{bmatrix} = \begin{bmatrix} A^2 A^d A^\pi & AA^d B D^\pi \\ 0 & D^2 D^d D^\pi \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}. \quad (18)$$

From ABD^π and DCA^π , we get

$$PP^\pi QW^\pi = \begin{bmatrix} 0 & AA^\pi B D^\pi \\ DD^\pi C A^\pi & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}.$$

Since $W^j W^\pi = 0$, $PP^\pi QW^\pi = 0$, $ABD^\pi = 0$ and $DCA^\pi = 0$, we have

$$\begin{aligned} \sum_{j=0}^{h-1} \left(\sum_{i=0}^{p-1} (Q^d)^{i+j+2} P^i \right) P^\pi QW^j W^\pi &= (Q^d)^2 P^\pi QW^\pi \\ &= \begin{bmatrix} (BC)^d & 0 \\ 0 & (CB)^d \end{bmatrix} \begin{bmatrix} A^\pi & 0 \\ 0 & D^\pi \end{bmatrix} \begin{bmatrix} 0 & B \\ C & 0 \end{bmatrix} \begin{bmatrix} A^\pi & 0 \\ 0 & D^\pi \end{bmatrix} \\ &= \begin{bmatrix} 0 & (BC)^d A^\pi B D^\pi \\ (CB)^d D^\pi C A^\pi & 0 \end{bmatrix} \\ &= \begin{bmatrix} 0 & (BC)^d B D^\pi \\ (CB)^d C A^\pi & 0 \end{bmatrix}. \end{aligned}$$

The conditions $ABD^\pi = 0$, $DCA^\pi = 0$ and $DD^d C = 0$ implies that $ABD^n C = 0$ and $DCBD^n C = 0$, for $n \geq 0$, observe that (16) and (18) yield

$$\begin{aligned} (P+Q)^{2i} P^\pi Q(W^d)^{2i+2} &= \begin{bmatrix} \sum_{i=0}^{n-1} B(CB)^i D^{2n-2i-1} C(A^d)^{2n+2} & \sum_{i=0}^{n-1} B(CB)^i D^{2n-2i-1} C Z_{2n+2} + \sum_{i=0}^n (BC)^i A^{2n-2i} A^\pi B(D^d)^{2n+2} \\ \sum_{i=0}^n (CB)^i D^{2n-2i} C(A^d)^{2n+2} & \sum_{i=0}^{n-1} (CB)^i D^{2n-2i} C Z_{2n+2} + \sum_{i=0}^{n-1} (CB)^i C A^{2n-2i-1} A^\pi B(D^d)^{2n+2} \end{bmatrix} \\ (P+Q)^{2n+1} P^\pi Q(W^d)^{2n+3} &= \begin{bmatrix} \sum_{i=0}^{n-1} B(CB)^i D^{2n-2i} C(A^d)^{2n+3} & \sum_{i=0}^n B(CB)^i D^{2n-2i} C Z_{2n+3} + \sum_{i=0}^n (BC)^i A^{2n+1-2i} A^\pi B(D^d)^{2n+3} \\ \sum_{i=0}^n (CB)^i D^{2n+1-2i} C(A^d)^{2n+3} & \sum_{i=0}^{n-1} (CB)^i D^{2n+1-2i} C Z_{2n+3} + \sum_{i=0}^n (CB)^i C A^{2n-2i} A^\pi B(D^d)^{2n+3} \end{bmatrix}. \end{aligned}$$

It follows that

$$\sum_{n=0}^{h-1} (P+Q)^n P^\pi Q(W^d)^{n+2} = \begin{bmatrix} \sum_{n=0}^{\left[\frac{h}{2}\right]-1} (P+Q)^{2n} P^\pi Q(W^d)^{2n+2} & \sum_{n=0}^{\left[\frac{h}{2}\right]-1} (P+Q)^{2n+1} P^\pi Q(W^d)^{2n+3} \end{bmatrix} = \begin{bmatrix} L & N \\ S & R \end{bmatrix},$$

where L, N, S and R are defined in (11).

Using the same way as (16), we have

$$\sum_{n=0}^{p-1} (Q^d)^{n+1} P^{n+1} = \sum_{n=0}^{\left[\frac{p}{2}\right]-1} (Q^d)^{2n+1} P^{2n+1} + \sum_{n=0}^{\left[\frac{p}{2}\right]-1} (Q^d)^{2n+2} P^{2n+2} = \begin{bmatrix} XA^2 & BYD \\ CXA & YD^2 \end{bmatrix}.$$

The condition $ABD^n C = 0$ implies that $AL = 0$, so we get

$$Q^\pi \sum_{j=0}^{h-1} (P+Q)^j P^\pi Q(W^d)^{j+2} - \sum_{j=0}^{h-1} \sum_{i=0}^{p-1} (Q^d)^{i+1} P^{i+1} (P+Q)^j P^\pi Q(W^d)^{j+2} = \begin{bmatrix} (BC)^\pi L - BYDS & (BC)^\pi N - XA^2N - BYDR \\ (CB)^\pi S - YD^2S & (CB)^\pi R - CXAN - YD^2R \end{bmatrix}.$$

The proof is finished.

The next result is a generalization of [9, Theorem 5].

Theorem 3.3 Let M be given by (1), if $DD^dC = 0$ and $BD^\pi = 0$. Then

$$M^d = \begin{bmatrix} A^d & -A^d BD^d + A^\pi \sum_{n=0}^{r-1} A^n B(D^d)^{n+2} \\ \sum_{n=0}^{s-1} D^n C(A^d)^{n+2} & D^d + \sum_{n=1}^{s-1} \sum_{i=1}^n D^{i-1} CA^{n-i} A^\pi B(D^d)^{n+2} + \sum_{n=0}^{s-1} D^n CX_{n+2} \end{bmatrix}$$

where

$$X_n = -\sum_{i=0}^{n-1} (A^d)^{i+1} B(D^d)^{n-i}, \quad n \geq 1. \quad (19)$$

Proof. We can split matrix M as $M = P + Q$, where

$$P = \begin{bmatrix} A^2 A^d & 0 \\ 0 & D^2 D^d \end{bmatrix}, \quad Q = \begin{bmatrix} AA^\pi & B \\ C & DD^\pi \end{bmatrix}, \quad P^d = \begin{bmatrix} A^d & 0 \\ 0 & D^d \end{bmatrix}, \quad P^\pi = \begin{bmatrix} A^\pi & 0 \\ 0 & D^\pi \end{bmatrix}.$$

From $DD^dC = 0$ and $BD^\pi = 0$, we have $PQP^\pi = 0$. Applying Theorem 2.1, we get

$$\begin{aligned} M^d &= W^d + \sum_{i=0}^{p-1} (Q^d)^{i+1} P^i P^\pi - \sum_{i=0}^{p-1} (Q^d)^{i+1} P^i P^\pi QW^d + \sum_{j=0}^{h-1} \left(\sum_{i=0}^{p-1} (Q^d)^{i+j+2} P^i \right) P^\pi QW^j W^\pi + Q^\pi \sum_{j=0}^{h-1} (P+Q)^j P^\pi Q(W^d)^{j+2} \\ &\quad - \sum_{j=0}^{h-1} \sum_{i=0}^{p-1} (Q^d)^{i+1} P^{i+1} (P+Q)^j P^\pi Q(W^d)^{j+2}, \end{aligned}$$

where $\text{ind}(P) = p$, $\text{ind}(Q) = q$, $\text{ind}(Q) = q$, $\max\{\text{ind}(P), \text{ind}(Q)\} \leq h \leq \text{ind}(P) + \text{ind}(Q)$, $\text{ind}(W) \leq \text{ind}(P) + \text{ind}(Q)$.

Since $DD^dC = 0$ and $BD^\pi = 0$, implies, for $n \geq 0$, $BD^n D^\pi C = BD^n C - BD^n DD^d C = BD^n C = 0$.

From $BD^\pi = 0$ and $BC = 0$, the matrix Q satisfies Lemma 3.3, by Lemma 3.2, we get

$$Q^d = \begin{bmatrix} (AA^\pi)^d & ((AA^\pi)^d)^2 B \\ \Sigma_0 & (DD^\pi)^d + \Sigma_1 B \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ \Sigma_0 & \Sigma_1 B \end{bmatrix},$$

where Σ_n is defined in (5), since $DD^dC = 0$, by Lemma 3.2, we have $\Sigma_n = 0$. Thus $Q^d = 0$ and $Q^\pi = I$. In this situation, we obtain

$$M^d = W^d + \sum_{j=0}^{h-1} (P+Q)^j P^\pi Q(W^d)^{j+2}.$$

Since $DD^dC = 0$, we have

$$W = PP^d(P+Q) = \begin{bmatrix} A^2 A^d & AA^d B \\ 0 & D^2 D^d \end{bmatrix}.$$

By Lemma 2.2 and Lemma 3.4, we get

$$W^d = \begin{bmatrix} A^d & X_1 \\ 0 & D^d \end{bmatrix}, (W^d)^n = \begin{bmatrix} (A^d)^n & X_n \\ 0 & (D^d)^n \end{bmatrix}.$$

where X_n is defined in (19). Clearly, for every integer $n \geq 0$,

$$P^\pi Q(W^d)^{j+2} = \begin{bmatrix} 0 & A^\pi B(D^d)^{j+2} \\ C(A^d)^{j+2} & CX_{j+2} \end{bmatrix}.$$

Since $DD^d C = 0$ and $BD^i C = 0$, we can prove, for every integer $n \geq 0$,

$$\sum_{j=0}^{h-1} (P+Q)^j P^\pi Q(W^d)^{j+2} = \begin{bmatrix} 0 & \sum_{n=0}^{r-1} A^\pi A^n B(D^d)^{n+2} \\ \sum_{n=0}^{s-1} D^n C(A^d)^{n+2} & \sum_{n=1}^{r-1} \sum_{i=1}^n D^{i-1} CA^{n-i} A^\pi B(D^d)^{n+2} + \sum_{n=0}^{s-1} D^n CX_{n+2} \end{bmatrix}$$

The proof is finished.

In the rest of the paper we will exploit Theorem 2.2 to obtain some representations of M^d under some weaker conditions. Firstly we will present the following result.

Theorem 3.4 Let M be given by (1), if $BD^\pi = 0$, $DD^d C = 0$ and $CAA^d = 0$. Then

$$M^d = \begin{bmatrix} A^d & X_1 + \sum_{n=0}^{r-1} A^\pi A^n B(D^d)^{n+2} \\ 0 & D^d + \sum_{n=1}^{r-1} \sum_{i=0}^{n-1} D^{n-i-1} CA^i B(D^d)^{n+2} \end{bmatrix},$$

where

$$X_n = -\sum_{i=0}^{n-1} (A^d)^{i+1} B(D^d)^{n-i}, n \geq 1. \quad (20)$$

Proof. We can split matrix M as $M = P + Q$, where

$$P = \begin{bmatrix} A & 0 \\ C & D \end{bmatrix}, Q = \begin{bmatrix} 0 & B \\ 0 & 0 \end{bmatrix}.$$

From $DD^d C = 0$ and $CAA^d = 0$, by Lemma 2.2, we get

$$P^d = \begin{bmatrix} A^d & 0 \\ 0 & D^d \end{bmatrix}, P^\pi = \begin{bmatrix} A^\pi & 0 \\ 0 & D^\pi \end{bmatrix}.$$

Since $DD^d C = 0$, we have, for every integer $n \geq 1$,

$$P^\pi P^n Q = \begin{bmatrix} 0 & A^\pi A^n B \\ 0 & \sum_{i=0}^{n-1} D^{n-i-1} CA^i B \end{bmatrix}.$$

Since $BD^\pi = 0$, we obtain $QP^\pi = 0$. Applying Theorem 2.2, we get

$$M^d = W^d + P^\pi \sum_{n=0}^{p-1} P^n Q(W^d)^{n+2}.$$

From $DD^dC = 0$ and $CAA^d = 0$, we have

$$W = PP^d(P+Q) = \begin{bmatrix} A^2A^d & AA^dB \\ DD^dC & D^2D^d \end{bmatrix} = \begin{bmatrix} A^2A^d & AA^dB \\ 0 & D^2D^d \end{bmatrix}.$$

Using the same way as Theorem 3.2, we obtain

$$W^d = \begin{bmatrix} A^d & X_1 \\ 0 & D^d \end{bmatrix}, (W^d)^n = \begin{bmatrix} (A^d)^n & X_n \\ 0 & (D^d)^n \end{bmatrix},$$

where X_n is defined in (20).

Hence, we get

$$P^\pi Q(W^d)^2 = \begin{bmatrix} 0 & A^\pi B \\ 0 & 0 \end{bmatrix} \begin{bmatrix} (A^d)^2 & X_2 \\ 0 & (D^d)^2 \end{bmatrix} = \begin{bmatrix} 0 & A^\pi B(D^d)^2 \\ 0 & 0 \end{bmatrix},$$

apparently

$$P^\pi \sum_{n=1}^{p-1} P^n Q(W^d)^{n+2} = \sum_{n=1}^{p-1} \begin{bmatrix} 0 & A^\pi A^n B \\ 0 & \sum_{i=0}^{n-1} D^{n-i-1} CA^i B \end{bmatrix} \begin{bmatrix} (A^d)^{n+2} & X_{n+2} \\ 0 & (D^d)^{n+2} \end{bmatrix} = \begin{bmatrix} 0 & \sum_{n=1}^{r-1} A^\pi A^n B(D^d)^{n+2} \\ 0 & \sum_{n=1}^{r-1} \sum_{i=0}^{n-1} D^{n-i-1} CA^i B(D^d)^{n+2} \end{bmatrix}.$$

The proof is finished.

The conditions $BD^\pi = 0$ and $CA^\pi = 0$ is less general than $ABD^\pi = 0$ and $DCA^\pi = 0$. The following expression for M^d is much more simpler than the Theorem 3.2.

Theorem 3.5 Let M be given by (1), if $BD^\pi = 0$, $DD^dC = 0$ and $CA^\pi = 0$. Then

$$M^d = \begin{bmatrix} A^d & X_1 + \sum_{n=0}^{r-1} A^\pi A^n B(D^d)^{n+2} \\ \sum_{n=0}^{s-1} D^n C(A^d)^{n+2} & D^d + \sum_{n=0}^{s-1} D^n CX_{n+2} \end{bmatrix},$$

where

$$X_n = -\sum_{i=0}^{n-1} (A^d)^{i+1} B(D^d)^{n-i}, n \geq 1. \quad (21)$$

Proof. We can represent P , P^d , P^π and Q as in (12). From $DD^dC = 0$, we have, for $n \geq 0$,

$$P^\pi P^n Q = \begin{bmatrix} 0 & A^\pi A^n B \\ D^n C & 0 \end{bmatrix}.$$

Since $BD^\pi = 0$ and $CA^\pi = 0$, we obtain $QP^\pi = 0$. Applying Theorem 2.2, we get

$$M^d = W^d + P^\pi \sum_{n=0}^{p-1} P^n Q(W^d)^{n+2}.$$

From $DD^dC = 0$, we have

$$W = PP^d(P+Q) = \begin{bmatrix} A^2A^d & AA^dB \\ DD^dC & D^2D^d \end{bmatrix} = \begin{bmatrix} A^2A^d & AA^dB \\ 0 & D^2D^d \end{bmatrix}.$$

Using the same way as Theorem 3.2, we have

$$W^d = \begin{bmatrix} A^d & X_1 \\ 0 & D^d \end{bmatrix}, (W^d)^n = \begin{bmatrix} (A^d)^n & X_n \\ 0 & (D^d)^n \end{bmatrix},$$

where X_n is defined in (21).

Hence, we get

$$\begin{aligned} & P^\pi \sum_{n=0}^{p-1} P^n Q (W^d)^{n+2} \\ &= \sum_{n=0}^{p-1} \begin{bmatrix} 0 & A^\pi A^n B \\ D^n C & 0 \end{bmatrix} \begin{bmatrix} (A^d)^{n+2} & X_{n+2} \\ 0 & (D^d)^{n+2} \end{bmatrix} \\ &= \begin{bmatrix} 0 & \sum_{n=0}^{p-1} A^\pi A^n B (D^d)^{n+2} \\ \sum_{n=0}^{s-1} D^n C (A^d)^{n+2} & \sum_{n=0}^{s-1} D^n C X_{n+2} \end{bmatrix}. \end{aligned}$$

The proof is finished.

Theorem 3.6. Let M be given by (1). If $BD^\pi = 0$, $A = BD^dC$ and $D^dCB = 0$. Then

$$M^d = \begin{bmatrix} B(D^d)^3C & B(D^d)^2 \\ (D^d)^2C + \sum_{n=1}^{s-1} D^{n-1}CB(D^d)^{n+3}C & D^d + \sum_{n=1}^{s-1} D^{n-1}CB(D^d)^{n+2} \end{bmatrix}.$$

Proof. We can split matrix M as $M = P + Q$, where

$$P = \begin{bmatrix} 0 & 0 \\ C & D \end{bmatrix}, Q = \begin{bmatrix} A & B \\ 0 & 0 \end{bmatrix}.$$

By Lemma 2.2, we have

$$P^d = \begin{bmatrix} 0 & 0 \\ (D^d)^2C & D^d \end{bmatrix}, P^\pi = \begin{bmatrix} I & 0 \\ -D^dC & D^\pi \end{bmatrix}.$$

From $BD^\pi = 0$, $A = BD^dC$ and $D^dCB = 0$, we have $AB = BD^dCB = 0$, $D^dCA = D^dCBD^dC = 0$.

From $D^dCA = 0$ and $D^dCB = 0$, we have, for $n \geq 1$,

$$P^\pi P^n Q = \begin{bmatrix} 0 & 0 \\ D^{n-1}CA & D^{n-1}CB \end{bmatrix}.$$

Since $BD^\pi = 0$ and $A = BD^dC$, we obtain $QP^\pi = 0$. Applying Theorem 2.2, we get

$$M^d = W^d + P^\pi \sum_{n=0}^{p-1} P^n Q (W^d)^{n+2}.$$

From $D^dCA = 0$ and $D^dCB = 0$, we have

$$\begin{aligned}
W &= PP^d(P+Q) \\
&= \begin{bmatrix} 0 & 0 \\ D^d C & DD^d \end{bmatrix} \begin{bmatrix} A & B \\ C & D \end{bmatrix} \\
&= \begin{bmatrix} 0 & 0 \\ D^d CA + DD^d C & D^d CB + D^2 D^d \end{bmatrix} \\
&= \begin{bmatrix} 0 & 0 \\ DD^d C & D^2 D^d \end{bmatrix},
\end{aligned}$$

By Lemma 2.2 and Lemma 3.2, we have

$$W^d = \begin{bmatrix} 0 & 0 \\ (D^d)^2 C & D^d \end{bmatrix}, (W^d)^n = \begin{bmatrix} 0 & 0 \\ (D^d)^{n+1} C & (D^d)^n \end{bmatrix}.$$

The conditions $D^d CA = 0$ and $D^d CB = 0$ implies that $P^\pi Q = Q$. Hence, we get

$$P^\pi Q(W^d)^2 = \begin{bmatrix} A & B \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & 0 \\ (D^d)^3 C & (D^d)^2 \end{bmatrix} = \begin{bmatrix} B(D^d)^3 C & B(D^d)^2 \\ 0 & 0 \end{bmatrix},$$

and

$$P^\pi \sum_{n=1}^{p-1} P^n Q(W^d)^{n+2} = \sum_{n=1}^{p-1} \begin{bmatrix} 0 & 0 \\ D^{n-1} CA & D^{n-1} CB \end{bmatrix} \begin{bmatrix} 0 & 0 \\ (D^d)^{n+3} C & (D^d)^{n+2} \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ \sum_{n=1}^{s-1} D^{n-1} CB(D^d)^{n+3} C & \sum_{n=1}^{s-1} D^{n-1} CB(D^d)^{n+2} \end{bmatrix}.$$

The proof is finished.

Theorem 3.6 has the following dual version.

Theorem 3.7 Let M be given by (1). If $CA^\pi = 0$, $D = CA^d B$ and $A^d BC = 0$. Then

$$M^d = \begin{bmatrix} A^d + \sum_{n=1}^{r-1} A^{n-1} BC(A^d)^{n+2} & (A^d)^2 B + \sum_{n=1}^{r-1} A^{n-1} BC(A^d)^{n+3} B \\ C(A^d)^2 & C(A^d)^3 B \end{bmatrix}.$$

Proof. Using the same way as Theorem 3.6, we can obtain the result.

Acknowledgements

The authors are very grateful to the reviewers for their suggestions. This research was supported by Guangxi Natural Science Foundation (No. 2016GXNSFAA380125).

References

- [1] M. P. Drazin, Pseudo-inverses in associative rings and semigroup, Amer. Math. Monthly, 65 (1958), 506-514.
- [2] R. E. Harte, Invertibility and Singularity, Dekker, 1988.
- [3] S. L. Campbell, The Drazin inverse and systems of second order linear differential equations, linear Multilinear Algebra, 14 (1983), 195-198.
- [4] S. L. Campbell, C. D. Meyer Jr., N. J. Rose, Applications of the Drazin inverse to linear systems of differential equations, SIAM J. Appl. Math., 31 (1976), 411-425.
- [5] S. L. Campbell, C. D. Meyer, Generalized Inverse of Linear Transformations, Pitman, London, 1979, Dover, New York, 1991.
- [6] N. Zhang, Y. Wei, Solving EP singular linear systems, Int. J. Comput. Math., 81 (2004), 1395-1405.
- [7] C. D. Meyer Jr., N. J. Rose, The index and the Drazin inverse of block triangular matrices, SIAM J. Appl. Math. 33 (1) (1977), 1-7.
- [8] R. E. Hartwig, J. M. Shoaf, Group inverse and Drazin inverse of bidiagonal and triangular toeplitz matrices, Austral. J. Math., 24 (A) (1977), 10-34.
- [9] G. Yan, X. Qin, X. Liu, Representations for the Drazin inverse of block matrices, Alabama Journal of Mathematics, 41 (2017): 1-10.
- [10] E. Dopazo, M. F. Martínez-Serrano, Further results on the representation of the Drazin inverse of a 2×2 block matrices, Linear Algebra Appl., 432 (2010), 1896-1904.
- [11] D. S. Cvetković-Ilić, A note on the representation for the Drazin inverse of 2×2 blockmatrices. Linear Algebra Appl., 429 (2008), 242-248.

- [12] D. Mosić, A note on the representations for the generalized Drazin inverse of block matrices, *Acta Mathematica Scientia*, 35B (2015): 1483–1491.
- [13] H. Yang, X. Liu, The Drazin inverse of the sum of two matrices and its applications, *J. Comput. Appl. Math.*, 235 (2011), 1412-1417.
- [14] L. Xia, B. Deng, The Drazin Inverse of the Sum of Two Matrices and its Applications, *Filomat*, 16 (2017), 5151–5158.
- [15] X. Liu, X. Qin, J. Benítez, New additive results for the generalized Drazin inverse in a Banach algebra, *Filomat*, 30 (2016), 2289-2294.
- [16] R. E. Hartwig, G. R. Wang, and Y. Wei, Some additive results on Drazin inverse, *Linear Algebra Appl.*, 322 (2001), 207-217.
- [17] M. Catral, D. D. Olesky, and P. Driessche, Block representations of the Drazin inverse of a bipartite block matrix, *Linear Algebra Appl.* 5 A (09) (2009), 98-107.
- [18] C. Deng, Y. Wei, A note on the Drazin inverse of an anti-triangular matrix, *Linear Algebra Appl.*, 431 (2009), 1910-1922.